

## Caridea postlarvae (Crustacea, Decapoda) in the estuary of the Jamapa River, Veracruz, SW Gulf of Mexico

## Postlarvas de Caridea (Crustacea, Decapoda) en el estuario del río Jamapa, Veracruz, SW golfo de México

Sergio Cházaro-Olvera<sup>1\*</sup>, Jesús Montoya Mendoza<sup>2</sup>, Ángel Morán-Silva<sup>1</sup>, Rafael Chávez-López<sup>1</sup>, Clever Ramírez-Isoba<sup>1</sup>

Recibido: 18 de marzo de 2024.

Aceptado: 17 de junio de 2024.

Publicado: diciembre de 2024.

### ABSTRACT

**Background.** In Veracruz, Mexico, 68 species of estuarine decapod crustaceans have been recorded. Of these, 11 belong to the infraorder Caridea, and 8 species have been recorded in freshwater systems. **Goals.** The present study analyzed the specific richness and abundance of carideans postlarvae and their relationship with the dynamics of environmental factors in the estuary of the Jamapa River, Veracruz, southwest of the Gulf of Mexico. **Methods.** The following environmental factors were measured: dissolved oxygen, temperature, total dissolved solids, pH, and salinity. The captures were made with light traps and shrimp bait. The generalized least squares model was used to determine the relationships between the environmental factors with the five months and five sampling sites. The generalized linear model was used to determine the relationships between the abundance of the postlarvae and the months, sites, and environmental factors. **Results.** A total of 8,649 carideans postlarvae were collected, of which 257 belonged to *Macrobrachium acanthurus*, 1,016 belonged to *M. olfersii*, and 7,376 belonged to *Potimirim mexicana*. The highest abundance of the three species was found at the site located on the dock of the Boca del Río Technological Institute, with 6,627 carideans. **Conclusion.** The abundances of postlarvae of *P. mexicana* and *M. olfersii* were within the values reported in other studies. However, the abundance of *M. acanthurus* was low, which may be related to the high values of total dissolved solids or overfishing. The highest density of postlarvae of carideans occurred in the rainy season in sites with *Thypha domingensis* vegetation.

**Keywords:** Atyidae, Caridean shrimps, diversity, estuarine system, Palaemonidae

<sup>1</sup> Laboratorio de Crustáceos. Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México. Av. de los Barrios No.1, Los Reyes Iztacala, Tlalnepantla, Estado de México, 54090. México

<sup>2</sup> Laboratorio de Investigación de Acuicultura Aplicada. Tecnológico Nacional de México/ Instituto Tecnológico de Boca del Río. Km 12, carretera Veracruz-Córdoba, Boca del Río, Veracruz, México

\*Corresponding author:

Sergio Cházaro-Olvera: schazaro@gmail.com

To quote as:

Cházaro-Olvera, S., J. Montoya Mendoza, Á. Morán-Silva, R. Chávez-López & C. Ramírez-Isoba. 2024. Caridea postlarvae (Crustacea, Decapoda) in the estuary of the Jamapa River, Veracruz, SW Gulf of Mexico. *Hidrobiología* 34 (3): 145-152.

DOI:10.24275/BSVD9975

### RESUMEN

**Antecedentes.** En Veracruz, México, se han registrado 68 especies de crustáceos decápodos de estuario. De ellas, 11 pertenecen al infraorden Caridea y 8 especies han sido registradas en sistemas de agua dulce. **Objetivos.** El presente estudio analizó la riqueza y abundancia específica de postlarvas de carideos y su relación con la dinámica de factores ambientales en el estuario del Río Jamapa, Veracruz, suroeste del Golfo de México. **Métodos.** Se midieron los siguientes factores ambientales: oxígeno disuelto, temperatura, sólidos disueltos totales, pH y salinidad. Las capturas se realizaron con trampas ligeras y cebo para camarones. Se utilizó el modelo de mínimos cuadrados generalizado para determinar las relaciones entre los factores ambientales con los cinco meses y los cinco sitios de muestreo. Se utilizó el modelo lineal generalizado para determinar las relaciones entre la abundancia de las postlarvas y los meses, sitios y factores ambientales. **Resultados.** Se recolectaron un total de 8,649 postlarvas de carideos, de las cuales 257 pertenecían a *Macrobrachium acanthurus*, 1,016 a *M. olfersii* y 7,376 a *Potimirim mexicana*. La mayor abundancia de las tres especies se encontró en el sitio ubicado en el muelle del Instituto Tecnológico de Boca del Río, con 6,627 carideos. **Conclusión.** Las abundancias de postlarvas de *P. mexicana* y *M. olfersii* estuvieron dentro de los valores reportados en otros estudios. Sin embargo, la abundancia de *M. acanthurus* fue baja, lo que puede estar relacionado con los altos valores de sólidos disueltos totales o la sobre pesca. La mayor densidad de postlarvas de carideos se presentó en la época de lluvias en sitios con vegetación de *Thypha domingensis*.

Palabras clave: Atyidae, Camarones carideos, diversidad, Palaemonidae, sistema estuarino

## INTRODUCTION

The infraorder Caridea is made up of 3,754 described species grouped into 45 families; among the most diverse are the families Atyidae, with 526 species, and Palaemonidae, with 1,090 species (WoRMS, 2024). In Veracruz, Mexico, 68 species of estuarine decapod crustaceans have been recorded. A total of 11 of these species belong to the infraorder Caridea. In freshwater systems, 38 species have been recorded, and 8 of those species are part of the infraorder (Álvarez et al., 2011).

The *Potimirim* genus from the Atyidae family and the *Machrobrachium* genus from the Palaemonidae family have been reported along the coastline and in the rivers of Veracruz (Alonso-Reyes et al., 2010; Cházaro-Olvera et al., 2021) and in estuaries and freshwater systems (Álvarez et al., 2011). During their development, the individuals from these families undergo four stages: zoea, postlarva, juvenile, and adult (Hernández-Vergara & Jiménez-Rojo, 2008). The postlarvae of these carideans present morpho-physiological adaptations to be able to carry out migrations between the river, estuary, and maritime waters near the coast (De Grave, 2008). This migration is classified as an active movement when the organisms can perform vertical migrations in the water column, or it can be a passive movement when the organisms are transferred by currents (Guerao, 1995). The transport of caridean larvae and postlarvae to estuaries is related with the river currents to the estuaries (Anger, 2013). Therefore, the passive dispersal of larvae and postlarvae may be a factor that determines the recruitment of carideans to estuarine systems, where they return to the river to freshwater conditions after completing development to the juvenile stage.

Postlarvae recruitment has only been documented by Cházaro-Olvera (1996), Cházaro-Olvera et al. (2007a), Cházaro-Olvera et al. (2007b), and Cházaro-Olvera et al. (2009). Thus, the present work will increase the knowledge about the dynamics of the transport of caridean postlarvae in the Jamapa River estuary, Veracruz, located southwest of the Gulf of Mexico. To do this, the present work aims to analyze the abundance and diversity of the organisms of this infraorder and determine their relationships with physicochemical factors.

## MATERIALS AND METHODS

### Study area

The Jamapa River basin is located between  $18^{\circ}45' - 19^{\circ}14'N$  and  $95^{\circ}56' - 97^{\circ}17'W$  (Fuentes-Mariles et al., 2014). The estuary discharges its waters in the Veracruzano Reef System National Park (PNSAV) (Liaño-Carrera et al., 2019). The estuary has a micro-tidal modulation of approximately 2.0 m, with biweekly synodic semidiurnal, diurnal, and lunisolar components (Salas-Monreal et al., 2019). The estuary also has a navigation channel in the southern part that generates important changes in its dynamics (Salas-Monreal et al., 2019). The shipping channel of the estuary produces strong currents of more than  $0.5 \text{ ms}^{-1}$  and a continuous exchange of brackish water with the ocean (Salas-Monreal et al., 2020) (Figure 1).

### Fieldwork

Specimens were collected at five sites. The first site was located to the south of the estuary, at the jetty of the Instituto Tecnológico de Boca del Río, Veracruz (ITBOCA). The second site was located to the north of the estuary at the jetty of the Instituto de Ciencias Marinas y Pesquerías of the Universidad Veracruzana (ICIMAP). The third site was located near

the estuarine inlet find of the Río Jamapa called Barco. The fourth and fifth sites (Venecia and Estero) were located to the southeast, in communication with the Mandinga Lagoon. The sampling campaign was carried out in September (i.e., the rainy season), November, January, March (i.e., the cold front season), and May (i.e., the dry season) of 2019. The biological material was collected over 12 h using a light trap (Cházaro-Olvera et al., 2018). The light trap was placed at the sampling sites at a depth of 0.5 m during the full moon phase because this lunar phase is when the effect of positive phototropism of zooplankton is maximized. The trap was placed at 20:00 h on the first sampling day and removed at 08:00 h the following day. Each sample was filtered through a 300 µm sieve and preserved in 0.5 L plastic bottles. Subsequently, the samples were fixed with 70% ethyl alcohol and labeled with information on the location, date, time, and type of sampling. The abiotic parameters of water temperature ( $^{\circ}\text{C}$ ), salinity, total dissolved solids (ppm), dissolved oxygen (mg L<sup>-1</sup>), and pH were measured in situ with a Hanna® HI 9828 multiparameter every month and at each site at the beginning and end of the sampling; later, the average and standard deviation were obtained.

### Laboratory work

The biological material was transported to the Crustacean Laboratory of the FES Iztacala. The material was reviewed, separated, and identified to the species level with the help of a stereoscopic microscope and an optical microscope following the criteria of Holthuis (1954) and Williams (1984).

### Statistical analysis

A generalized least squares (GLS) model was used to determine the relationships between the environmental factors, the five months, and the five sampling sites (Zuur et al., 2007). After finding statistically significant differences between the means of the environmental factors of the months and sampling sites, Tukey's post hoc test was applied (Sokal & Rohlf, 1995). A generalized linear model (GLM) was used to determine the relationships between the abundance of caridean species in the postlarva stage with respect to sites, months, and environmental factors. A Poisson logarithmic linear (per counts) model was used, considering each species' abundance as dependent variable, the months and sampling sites as fixed factors and the environmental factors were considered as independent variables. A type III analysis was performed, and the chi-square statistic was obtained using the Wald model. Previously, the values of the environmental factors were transformed to arcsine and the abundance values of the species to  $\log(n+1)$  (Zuur et al., 2007). The GLS and GLM analyses were performed using SPSS Statistics 25 software of IBM corporation.

## RESULTS

### Environmental parameters

The concentration of dissolved oxygen had a range of  $3.65 \pm 0.65 \text{ mg L}^{-1}$  in November to  $6.65 \pm 0.04 \text{ mg L}^{-1}$  in May (Table 1). The GLS test showed a statistically significant relationship between the dissolved oxygen and the sampling months ( $r = 0.74$ ;  $P < 0.001$ ; Table 2). Statistically significant differences were not found between the dissolved oxygen concentrations of the sampling sites ( $P < 0.05$ ). However, statistically significant differences were found between the sampling months ( $P < 0.05$ ). After applying Tukey's post hoc test, only significant differences between May and September were found ( $P < 0.05$ ).




Figure 1. Locations of sampling sites in the estuary of the Río Jamapa, Boca del Río, Veracruz, México. ITBOCA: Instituto Tecnológico de Boca del Río; ICIMAP: Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana.

The temperature ranged from  $21.66 \pm 0.04^\circ\text{C}$  in January to  $31.64 \pm 0.11^\circ\text{C}$  in May (Table 1). The test GLS showed that there was a positive significant relationship between the temperature with the sites and months of sampling ( $r = 0.73$ ;  $P < 0.05$ ) (Table 2). Significant differences were observed among the sampling months ( $P < 0.001$ ) (Table 2). When applying Tukey's test, statistically significant differences between May and January, March and Nov, and January and September were found ( $P < 0.05$ ).

Salinity ranged from  $0.97 \pm 0.06$  in September to  $35.65 \pm 0.01$  in May (Table 1). Salinity showed a high correlation with the sites and sampling months ( $r = 0.91$ ;  $P < 0.001$ ) (Table 2). Significant differences were found between sites and between months ( $P < 0.001$ ). After applying Tukey's post hoc test, significant statistical differences were found between the ITBOCA, ICIMAP, and Barco sites and the Esteros and Venecia sampling sites ( $P < 0.05$ ). With respect to months, May differed from the other five sampling months. January was significantly different from September ( $P < 0.05$ ).

The pH ranged from  $7.03 \pm 0.46$  in November to  $7.85 \pm 0.06$  in January (Table 1). The GLS test showed a significant positive relationship between pH and the sites and months of sampling ( $r = 0.79$ ;  $P < 0.001$ ). Significant differences were found only between the months sampled ( $P < 0.001$ ) (Table 2). Tukey's test showed that there were only significant differences between January and September ( $P < 0.05$ ).

The range of total dissolved solids was  $10.95 \pm 1.05$  ppm in May and  $5,842 \pm 856$  ppm in March (Table 1). The relationship between total dissolved solids and the sites and months of sampling was positive ( $r = 0.62$ ) and significant ( $P < 0.05$ ). Only the sites presented significant differences ( $P < 0.05$ ) (Table 2). Tukey's test only showed significant differences between the ICIMAP and Venecia sites ( $P < 0.05$ ).

#### Abundance and specific richness

In total, 8,649 caridean postlarvae were collected, of which 257 belonged to *Macrobrachium acanthurus* (Weigmann, 1836), 1,016 belonged to *Macrobrachium olfersii* (Weigmann, 1836), and 7,376 belonged to *Potimirim mexicana* (De Saussure, 1857). The greatest abundance of *M. acanthurus*, *M. olfersii*, and *P. mexicana* was found at the ITBOCA site with 6,627 postlarvae, and the lowest abundance was found at Venecia with 361 postlarvae. According to the collection months, the highest abundance of the three species was found in September with 5,030 postlarvae, and the lowest abundance was found in March with only 40 postlarvae (Table 3).

When the GLM analysis was applied to the abundance of postlarvae of *P. mexicana*, it was observed that the variation was related to the five environmental factors ( $P < 0.05$ ). The abundance of *M. acanthurus* postlarvae was related to the dissolved oxygen and total dissolved solids ( $P < 0.05$ ). The abundance of *M. olfersii* postlarvae was related to dissolved oxygen, temperature, total dissolved solids, and salinity ( $P < 0.05$ ; Table 4).

## DISCUSSION

In the study area, dissolved oxygen values have been reported to range from 5.63 mg L<sup>-1</sup> in the dry season to 5.55 mg L<sup>-1</sup> in the cold front season and 5.35 mgL<sup>-1</sup> in the rainy season (Castañeda-Chávez *et al.*, 2017). These values are consistent with the dissolved oxygen values found in this study. In September, November, and January (i.e., the end of the rainy season and the cold front season), the concentration of dissolved oxygen decreased. This may be because this area has been classified as an urban estuary, so there are discharges of wastewater to

the river (Castañeda-Chávez *et al.*, 2017; Salas-Monreal *et al.*, 2020). Furthermore, the high quantity of organic matter transported in the rainy season by the river's own water may decrease the concentration of dissolved oxygen. The dissolved oxygen concentration enables the species of *Macrobrachium* to be present in the estuary. In this regard, Urbano *et al.* (2010) mentioned that values around  $5.30 \pm 2.15$  mgL<sup>-1</sup> are within the normal range for the cultivation of postlarvae of most river prawn species. Likewise, Mires (1983) pointed out that the dissolved oxygen concentration suitable for the survival of shrimp postlarvae of the species *M. rosenbergii* (De Man, 1879) is 2.5–8.4 mgL<sup>-1</sup>.

Table 1. Environmental factors of the Jamapa River estuary, Boca del Río, Veracruz. ITBOCA: Instituto Tecnológico de Boca del Río; ICIMAP: Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana; SD, standard deviation. DO: Dissolved oxygen, Sal: salinity, Temp: temperature, Tds: total dissolved solids.

Month	Sampling site	DO (mgL <sup>-1</sup> )	pH	Temp (°C)	Sal (PSU)	Tds (ppm)
September	ITBOCA	5.28 ± 0.06	7.18 ± 0.01	28.47 ± 0.65	0.97 ± 0.06	890 ± 50
	ICIMAP	3.88 ± 0.96	7.29 ± 0.09	30.05 ± 1.02	1.07 ± 0.23	904 ± 31
	Barco	4.02 ± 0.02	7.18 ± 0.04	29.51 ± 0.58	2.7 ± 0.75	1874 ± 106
	Venecia	4.85 ± 0.22	7.13 ± 0.01	29.48 ± 0.66	8.31 ± 0.93	258 ± 44
	Esteros	4.20 ± 0.14	7.19 ± 0.06	20.11 ± 0.17	14.29 ± 1.46	1054 ± 78
	Average ± SD	4.44 ± 0.06	7.19 ± 0.06	27.52 ± 4.18	5.47 ± 5.77	996 ± 578
November	ITBOCA	5.38 ± 0.26	7.75 ± 0.26	22.64 ± 0.66	6.16 ± 0.52	4688 ± 1648
	ICIMAP	3.99 ± 0.46	7.03 ± 0.46	24.32 ± 0.54	6.09 ± 1.13	4493 ± 363
	Barco	3.65 ± 0.65	7.16 ± 0.65	27.64 ± 0.11	5.22 ± 0.59	4226 ± 147
	Venecia	4.92 ± 0.1	7.50 ± 0.1	26.09 ± 0.37	22.04 ± 0.46	18.13 ± 1.18
	Esteros	4.96 ± 0.0	7.2 ± 0.07	26.45 ± 0.87	22.39 ± 1.12	18.16 ± 1.34
	Average ± SD	4.58 ± 0.72	7.33 ± 0.29	25.48 ± 1.97	12.38 ± 8.99	2689 ± 2244
January	ITBOCA	6.01 ± 0.01	7.44 ± 0.01	23.08 ± 0.78	7.04 ± 1.20	5006 ± 626
	ICIMAP	5.11 ± 0.15	7.65 ± 0.03	21.66 ± 0.04	7.46 ± 1.65	5091 ± 617
	Barco	4.11 ± 0.14	7.34 ± 0.02	22.6 ± 0.07	12.89 ± 0.95	10.95 ± 1.05
	Venecia	4.31 ± 0.08	7.85 ± 0.06	22.13 ± 0.71	34.03 ± 0.99	25.97 ± 0.47
	Esteros	4.75 ± 0.21	7.34 ± 0.02	22.38 ± 0.87	33.65 ± 0.36	25.53 ± 0.24
	Average ± SD	4.85 ± 0.75	7.52 ± 0.22	22.37 ± 0.53	19.01 ± 13.73	2032 ± 2754
March	ITBOCA	6.11 ± 0.14	7.16 ± 0.01	24.61 ± 0.57	6.14 ± 0.26	4913 ± 490
	ICIMAP	5.44 ± 0.32	7.24 ± 0.04	24.42 ± 0.21	8.02 ± 0.86	5842 ± 856
	Barco	5.15 ± 0.05	7.18 ± 0.01	24.81 ± 0.28	17.99 ± 2.33	14.36 ± 1.42
	Venecia	5.07 ± 0.06	7.19 ± 0.02	22.97 ± 0.51	33.93 ± 0.81	27.96 ± 2.37
	Esteros	6.19 ± 0.03	7.21 ± 0.04	22.66 ± 0.14	34.99 ± 0.43	27.86 ± 1.6
	Average ± SD	5.59 ± 0.53	7.2 ± 0.03	23.89 ± 0.99	20.21 ± 13.77	2165 ± 2950
May	ITBOCA	4.87 ± 0.21	7.62 ± 0.10	31.64 ± 0.11	33.37 ± 0.27	26.04 ± 0.83
	ICIMAP	6.65 ± 0.04	7.34 ± 0.01	30.45 ± 0.17	33.84 ± 0.22	26.89 ± 0.18
	Barco	6.06 ± 0.04	7.53 ± 0.04	28.91 ± 0.16	35.65 ± 0.01	27.57 ± 0.76
	Venecia	5.7 ± 0.37	7.56 ± 0.37	28.81 ± 0.32	35.39 ± 0.54	26.89 ± 0.18
	Esteros	5.43 ± 0.27	7.44 ± 0.27	28.92 ± 0.23	35.57 ± 0.04	27.13 ± 0.28
	Average ± SD	5.74 ± 0.67	7.49 ± 0.11	29.74 ± 1.26	34.76 ± 1.08	26.9 ± 0.56

Table 2. Generalized least squares model (GLS) for environmental factors registered in the inlet of the River Jamapa, Boca del Río, Veracruz during 2018 and 2019. df: degrees of freedom, Do: Dissolved oxygen, F: statistic in ANOVA (analysis of variance), Temp: temperature, Tds: total dissolved solids, P: probability level.

Origin	Do		pH		Temp °C		Tds		Sal	
	df	F	P	F	P	F	P	F	P	F
Corrected model	8	5.79	0.001	7.44	<0.001	5.39	0.002	3.19	0.023	18.74
Intersection	1	360.44	<0.001	11810.17	<0.001	3172.72	<0.001	23.01	<0.001	286.41
Site	4	1.98	0.15	3.27	0.04	1.66	0.21	4.31	0.015	14.17
Month	4	9.62	<0.001	11.6	<0.001	9.13	<0.001	2.09	0.13	23.29
Error	16									
Total	25									
Correlation coefficient (r)		0.74		0.79		0.73		0.62		0.91

In this region of the Gulf of Mexico, in the cold front season, Jasso-Montoya (2012), Avendaño-Alvarez (2013), Contreras-Espinoza (2016), and Castañeda-Chávez *et al.* (2017) reported that the temperature decreases to between 23°C and 24°C. In the study zone, Cházaro-Olvera *et al.* (2022) registered a temperature of 25.11°C ± 0.12 °C in ITBOCA in the cold front season, while Contreras-Espinoza (2016) mentioned that the Jamapa River temperature was 25°C in the cold front season and 29.4°C in the rainy season. Therefore, the temperature values in the Jamapa River estuary are consistent with the behavior of the region's climatic seasons (Zavala-Hidalgo *et al.*, 2006). We consider it important to use the Mexican regulations (NOM-001-SEMARNAT-2021) to compare the values obtained in this study. The temperature registered in this work does not exceed the maximum permissible limit of 35°C defined by the official Mexican standard. Existing research shows that the growth of the different stages of development of *M. americanum* (Spence Bate 1868) is optimal at a temperature of 26–29°C (López-Uriostegui *et al.*, 2020). Cházaro-Olvera *et al.* (2022) recollected postlarvae of *M. acanthurus* and *M. olfersii* in these temperature values.

In the Jamapa River estuary, a range of salinity values have been recorded from 2 psu in the rainy season to 21 psu in the dry season

(Aké-Castillo *et al.*, 2016; González-Vázquez *et al.*, 2019). In the present work, a wide variation of salinity was also observed. Some river prawn species, such as *M. americanum*, develop adequately between 3 and 15 psu (Chung, 2001; López-Uriostegui *et al.*, 2020). This implies that the *Macrobrachium* river prawns require brackish water in their post-larvae development, while juveniles and adults prefer low-salinity water or freshwater (Graziani *et al.*, 1995).

Lorán-Nuñez (2013) reported an abundance of 69–120 *M. acanthurus* postlarvae and juveniles in the lower basin of the Papaloapan River. This range is similar to the abundance identified in the present study in the estuary of the Jamapa River, where the abundance was 18–164 in the months of capture. The highest abundance was found in September (i.e., the rainy season) at the ITBOCA sampling site, which was characterized by the presence of *Thypha domingensis* Persoon 1807 vegetation. This finding is consistent with what was reported by Lorán-Nuñez (2013), who found the highest values of abundance in the rainy season. It is important to mention that in a study carried out at another latitude on the banks of the Iguaçu River in São Paulo, Brazil, the abundance of *M. olfersii* was 23,818 postlarvae, juveniles and adults (Ribeiro *et al.*, 2020).

Table 3. Carideans of the Jamapa River estuary. Abundances of five places and five months.

Species/Community Factor	Site					
	ITBOCA	ICIMAP	Barco	Venezia	Estero	Total
<i>Macrobrachium acanthurus</i> (Weigmann, 1836)	223	23	0	0	11	257
<i>Macrobrachium olfersii</i> (Weigmann, 1836)	658	181	62	50	65	1016
<i>Potimirim mexicana</i> (De Saussure, 1857)	5746	188	805	311	326	7376
Total	6627	392	867	361	402	8649
Month						
Species/Community Factor	September	November	January	March	May	Total
	164	34	20	18	21	257
<i>Macrobrachium acanthurus</i> (Weigmann, 1836)	609	122	98	10	177	1016
<i>Potimirim mexicana</i> (De Saussure, 1857)	4257	533	2436	12	138	7376
Total	5030	689	2554	40	336	8649

In the Jamapa River basin, pH values ranging from 6 to 9 have been recorded (Houbron, 2010; SEMARNAT, 2002). The pH values recorded in this study were neutral to slightly alkaline and were among the values established by the official Mexican standard (NOM-127-SSA1-1994). In the Jamapa River estuary, the pH values provide a buffer effect to the water, avoiding acidification (Bates, 1973). Pretto (1988) found that for good development of postlarvae and juveniles of *M. rosenbergii* shrimp, pH must range between 7 and 9.

The total dissolved solids were highest in the cold front and rainy seasons. Cházaro-Olvera *et al.* (2022) also found high values of total dissolved solids (732–1,443 ppm) in the cold front season. In the Jamapa River estuary, total dissolved solids in some months and sites exceeded the maximum permissible limit of 1,000 ppm, which was established by the official Mexican standard (NOM-001-SEMARNAT-2021 and NOM-127-SSA1-1994 for drinking water). These concentrations of dissolved solids are due to the transport from the lower basin of the Jamapa River to the estuary (Aragón-López *et al.*, 2017), which can affect the respiration process of carideans.

When analyzing the abundance of caridean postlarvae in this study, *P. mexicana* was the most abundant species. This finding is consistent with what was reported in the lower basin of the Papaloapan River by Miranda-Vidal *et al.* (2016), who collected 4,587 crustaceans, with *P. mexicana* being the dominant species at 34% in the dry season; *Macrobrachium* sp. and *P. mexicana* represented 59% and 30% of the abundance, respectively, and were the dominant taxa in the rainy season.

Specific richness is closely related to the dynamics of salinity in estuaries. For example, Barba *et al.* (2005) reported nine species of

carideans in Laguna Madre, Tamaulipas, where two of them were associated with submerged vegetation or littoral vegetation. On the other hand, the authors reported 11 species of carideans in Laguna de Terminos, Campeche, where five species were associated with submerged vegetation. In each of the lagoons the authors found at least one dominant species in the taxocene, which is consistent with the present study where the postlarvae of *P. mexicana* were dominant. In the Papaloapan River, Miranda-Vidal *et al.* (2016) also found that *P. mexicana* in the dry season and *Macrobrachium* sp. in the rainy season, respectively, were the dominant species.

In conclusion, the values of the environmental factors were related with the sites and months of sampling. The abundance values for *P. mexicana*, *M. acanthurus*, and *M. olfersii* were within the interval of the values obtained by other authors. The abundances of *P. mexicana* and *M. olfersii* were within the values reported in other studies. However, the abundance of *M. acanthurus* was relatively low, which may be related to the high values of total dissolved solids or overfishing. Finally, the highest density of caridean postlarvae occurred in the rainy season in sites with *T. domingensis* vegetation.

## ACKNOWLEDGMENTS

We appreciate the support received from the Boca del Río Veracruz Technological Institute for this study. We thank the reviewers for the work done and the time dedicated to reviewing this study.

## Conflict of interests

The authors declare that there is no conflict of interest.

Table 4. Relationship with GLM of abundance with respect to months, places and environmental factors. B, regression coefficient; Wald's  $\chi^2$  to determine if the explanatory variable of the model is significant; \* significant relationship.

Source	<i>M. acanthurus</i>			<i>M. olfersii</i>			<i>P. mexicana</i>		
	B	$\chi^2$ Wald	p	B	$\chi^2$ Wald	p	B	$\chi^2$ Wald	p
Intersection	7.97	395.96	< 0.001*	3.35	8.64	0.003*	6.17	53.05	< 0.001*
Barco	2.99	221.21	< 0.001*	-2.50	18.24	< 0.001*	1.69	21.42	< 0.001*
Estero	-1.28	14.67	< 0.001*	-1.89	7.97	0.005*	0.66	2.37	0.24
ICIMAP	9.34	583.01	< 0.001*	-0.09	0.01	0.909	9.91	312.92	< 0.001*
ITBOCA	1.75	141.34	< 0.001*	-0.81	5.51	0.019*	1.95	51.88	< 0.001*
Venecia	0 <sup>a</sup>			0 <sup>a</sup>			0 <sup>a</sup>		
Enero	2.29	460.60	< 0.001*	-0.47	1.07	0.300	1.23	45.95	< 0.001*
March	8.94	783.37	< 0.001*	2.71	32.18	< 0.001*	5.88	194.10	< 0.001*
May	2.54	314.07	< 0.001*	1.61	18.01	< 0.001*	2.19	71.54	< 0.001*
November	4.96	1068.88	< 0.001*	2.01	26.05	< 0.001*	3.27	154.83	< 0.001*
September	0 <sup>a</sup>			0 <sup>a</sup>			0 <sup>a</sup>		
Dissolved Oxygen mgL <sup>-1</sup>	0.002	54.27	< 0.001*	0.002	7.78	0.005*	0.002	15.01	< 0.001*
Ph	0.001	9.15	0.002*	0.001	0.21	0.654	0.001	0.69	0.405
Temperature °C	-0.002	211.96	< 0.001*	0.001	0.75	0.386	-0.001	94.20	< 0.001*
Total dissolved solids ppm	-0.001	1543.34	< 0.001*	0.001	48.75	< 0.001*	-0.001	297.61	< 0.001*
Salinity psu	-0.004	751.45	< 0.001*	0.001	3.52	0.061	-0.003	351.28	< 0.001*

a. zero because this parameter is redundant.

## REFERENCES

- Aké-Castillo, J. A., C. F. Rodríguez-Gómez, H. Perales-Valdina & R. Sanay-González. 2016. Florecimiento de *Heterocapsa rotundata* (Dinophyta) en el estuario río Jamapa, Veracruz: 323-333. In: García-Mendoza E, Quijano-Scheggia SI, Olivos-Ortiz A, Núñez-Vázquez EJ (eds.). Florecimientos Algales Nocivos en México. Ensenada, México. CICESE 438 p.
- ALONSO-REYES, P., BORTOLINI-ROSALES, J. L. AND ÁLVAREZ, F. (2010). Análisis discriminante aplicado a los grupos sexuales de *Potimirim mexicana*, camarón hermafrodita protándrico. *Revista Mexicana de Biodiversidad* 81, 187–192.
- ÁLVAREZ, F., J.L. VILLALOBOS & S. CHÁZARO-OLVERA. 2011. Camarones y cangrejos dulceacuícolas y marinos (Crustacea: Decapoda) A. Cruz-Angón (Ed.), La biodiversidad en Veracruz, estudio de estado, CONABIO, México D.F.: 287-294.
- Anger, K. (2013). Neotropical Macrobrachium (Caridea: Palaemonidae): on the biology, origin, and radiation of freshwater-invading shrimp. *Journal of Crustacean Biology* 33(2), 151–183. doi.org/10.1163/1937240x-00002124
- ARAGÓN-LÓPEZ, P., M. R. CASTAÑEDA-CHÁVEZ, A. GRANADOS-BARBA, D. SALAS-MONREAL & C. LANDEROS-SÁNCHEZ. 2017. Lindane and Endosulfan Sulfate Isomers in *Crassostrea virginica* (Gmelin, 1791) Oyster Populations in Lagoon Systems from Central Gulf of Mexico. *International Journal of Agriculture Environment and Biotechnology* 2 (4): 1579-1585. DOI: 10.22161/ijeab/2.4.17
- AVENDAÑO-ÁLVAREZ, J. O. 2013. Variación hidrológica intermensual del Sistema Arrecifal Veracruzano. Tesis de Maestría, Universidad Veracruzana, Veracruz, México. 97 p.
- BARBA, E., A. RAZ-GUZMAN & A. J. SÁNCHEZ. 2005. Distribution patterns of estuarine caridean shrimps in the southwestern Gulf of Mexico. *Crustaceana* 78: 709-726. DOI: 10.1163/156854005774353502
- BATES, R. G. 1973. Determination of pH: theory and practice. John Wiley & Sons, New York. 479 p.
- CASTAÑEDA-CHÁVEZ, M. R., A. C. SOSA-VILLALOBOS, I. A. AMARO-ESPEJO, I. GALAVIZ-VILLA & F. LANGO-REYNOSO. 2017. Eutrophication in the lower coastal basin of the Jamapa River in Veracruz, Mexico. *International Journal Resesearch Granthaalayah* 5 (12): 206-216. DOI: 10.29121/granthaalayah.v5.i12.2017.495
- CHÁZARO-OLVERA, S. 1996. Descripción de megalopas de las especies *Callinectes sapidus* Rathbun, *C. similis* William, *C. gracilis* (Sausse) de la boca de la Laguna Camaronera, Alvarado, Ver. Tesis de Maestría en Ciencias. Facultad de Ciencias, UNAM, México. 113 p.
- CHÁZARO-OLVERA, S., A. ROCHA-RAMÍREZ, A. RAMÍREZ-ROJAS, H. VÁZQUEZ-LÓPEZ, R. CHÁVEZ-LÓPEZ. 2007a. Recruitment of *Callinectes sapidus* Rathbun 1896 megalopae from three southwestern Gulf of Mexico lagoon-system inlets. *International Journal Zoological Research* 3 (3): 145-156. DOI: 10.3923/ijzr.2007.145.156
- CHÁZARO-OLVERA, S., A. ROCHA-RAMÍREZ & P. ARELLANO-RODRARTE. 2007b. Transport of *Pachygrapsus gracilis* (de Saussure, 1858) megalopae from a lagoon system inlet in the southwestern Gulf of Mexico. *Crustaceana* 80 (8) :955-968. DOI: 10.1163/156854007781681337
- CHÁZARO-OLVERA, S., I. WINFIELD & V. CORIA-OLVERA. 2009. Transport of *Faxonius aztecus* postlarvae in three lagoon-system inlets in the southwestern Gulf of Mexico. *Crustaceana* 82 (4): 425-437. DOI:10.1163/156854008X400612
- CHÁZARO-OLVERA, S., R. ROBLES, J. MONTOYA-MENDOZA & J. A. HERRERA-LÓPEZ. 2018. Intraspecific variation in megalopae of *Clibanarius antillensis* (Anomura, Diogenidae) among western Atlantic populations. *Nauplius* 26: e2018031. DOI: 10.1590/2358-2936e2018031
- CHÁZARO-OLVERA, S., A. ARIAS-MARTÍNEZ, J. MONTOYA-MENDOZA, Á. MORÁN-SILVA, R. CHÁVEZ-LÓPEZ, H. VÁZQUEZ-LÓPEZ, & A. DEL C. RODRÍGUEZ-VARELA. 2021. Larvae and macro-crustaceans along the coastline of the Parque Nacional Sistema Arrecifal Veracruzano, SW Gulf of Mexico. *Nauplius* 29. DOI: 10.1590/2358-2936e2021047
- CHÁZARO-OLVERA, S., MONTOYA-MENDOZA, J., CASTAÑEDA-CHÁVEZ, M. DEL R., LANGO-REYNOSO, F., & CHAPARRO-MEDINA, V. 2022. Acute toxicity of the herbicide karmex (diuron) in *Macrobrachium acanthurus* and *M. olfersii* prawns. *Revista Internacional de Contaminación Ambiental* 38: 381–390. DOI: 10.20937/rica.5461
- CHUNG, K. S. 2001. Adaptabilidad ecofisiológica de organismos acuáticos tropicales a cambios de salinidad. *Revista de Biología Tropical* 49 (1): 9-13.
- CONTRERAS-ESPINOZA M. 2016. Variación espaciotemporal de la estructura comunitaria del zooplancton y su relación con las variables hidrográficas en la desembocadura del río Jamapa, Veracruz, en dos temporadas climáticas (nortes y lluvias). Tesis Maestría en Ecología y Pesquerías, Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana, Veracruz, México. 90 p.
- RIBEIRO, C.C., LOPES, V.H., AND BERTINI, G. 2020. Abundance and spatio-temporal distribution of the amphidromous shrimp *Macrobrachium olfersii* (Caridea: Palaemonidae) along the Ribeira de Iguaape River (São Paulo, Brazil). *Nauplius* 28: e2020017. DOI: 10.1590/2358-2936e2020017
- DE GRAVE, S., CAI, Y., AND ANKER, ARTHUR. 2008. "Global diversity of shrimps (Crustacea: Decapoda: Caridea) in freshwater." *Hydrobiologia*, 287–293. 595. DOI: 10.1007/s10750-007-9024-2
- FUENTES-MARILES, O. A., V. FRANCO, F. LUNA-CRUZ, L. VÉLEZ-MORALES & H. L. MORALES-RODRÍGUEZ. 2014. Caracterización fluvial e hidráulica de las inundaciones en México convenio CNA-SGT-GASIR-09/2014 organismo de cuenca X Golfo centro ciudad de Veracruz, Veracruz ríos Jamapa y Cotaxtla. Comisión Nacional del Agua e Instituto de Ingeniería, UNAM. Mexico. 83 p.
- GONZÁLEZ-VÁZQUEZ, J. A., E. HERNÁNDEZ-VIVAR, C. ROJAS-SERNA & J. DEL VALLE-MORALES 2019. Diagnosis of water circulation in an estuary: A case study of the Jamapa River and the Mandinga lagoons, Veracruz, Mexico. *Ciencias Marinas* 45 (1): 1-16. DOI:10.7773/cm.v45i1.2923
- GUERAO, G. 1995. Locomotor activity patterns and feeding habits in the prawn *Palaemon xiphias* (Crustacea: Decapoda: Palaemonidae) in Alfacas Bay, Ebro Delta (northwest Mediterranean). *Marine Biology* 122(1):115-119.

- GRAZIANI, C. A., M. DE DONATO, K. S. CHUNG. 1995. Salinidades óptimas en larvas y postlarvas de *Macrobrachium carcinus* (L.) (Decapoda: Palaemonidae). *Boletín del Instituto de Oceanografía de Venezuela* 34: 33-40.
- HERNÁNDEZ-VERGARA, M. P. & S. JIMÉNEZ-ROJO. 2008. Desarrollo larval y supervivencia de *Atya scabra* (Crustacea: Decapoda: Atyidae), a diferentes salinidades de cultivo. *Ciencia Pesquera* 16:15-22.
- HOLTHUIS, L. B. 1954. On a collection of Decapod Crustacea from the Republic of El Salvador (Central America). *Zoologische Verhandelingen* 23 (1): 1-43.
- HOUBRON, E. 2010. Calidad del agua. In: Atlas del patrimonio natural, histórico y cultural de Veracruz; Enrique Florescano, Juan Ortíz Escamilla, coordinadores. México: Gobierno del Estado de Veracruz: Comisión del Estado de Veracruz para la Conmemoración de la Independencia Nacional y la Revolución Mexicana: *Universidad Veracruzana* 1:147-159.
- JASSO-MONTOYA, J. 2012. Variación de los parámetros oceanográficos alrededor del Arrecife Verde en el Parque Nacional Sistema Arrecifal Veracruzano (Golfo de México Occidental). Tesis de Maestría en ciencias, Universidad Veracruzana, Boca del Río, Veracruz, México. 71 p.
- LIAÑO-CARRERA, F., T. CAMARENA-LUHRS, A. GÓMEZ-BARREIRO, F. J. MARTOS-FERNÁNDEZ, J. I. RAMÍREZ-MACIAS, D. SALAS-MONREAL. 2019. New coral reef structures in a tropical coral reef system. *Latin american journal of aquatic research* 47 (2): 270-281. DOI: 10.3856/vol47-issue2-fu-lltext-7
- LÓPEZ-URIESTEGUI, F., J. T. PONCE-PALAOFOX, F. LANGO-REYNOSO, M. R. CASTAÑEDA-CHÁVEZ, I. GALAVÍZ-VILLA, S. CASTILLO-VARGASMACHUCA & A. RUIZ-LUNA. 2020. A Response-Surface Analysis of Temperature Salinity on Growth and Survival of the Caridean Shrimp, *Macrobrachium americanum* (Bate, 1868). *Pakistan Journal of Zoology* 52 (5): 1903-1909 DOI: 10.17582/journal.pjz/20180523170525
- LORÁN-NÚÑEZ, R. M. 2013. Aspectos poblacionales del langostino *Macrobrachium acanthurus* (Wiegmann, 1836) en algunos lugares de la cuenca baja del río Papaloapan, Veracruz, México. Tesis de Maestría en Ciencias (Biología), Facultad de Ciencias, Universidad Nacional Autónoma de México. México 71 p.
- MIRANDA-VIDAL, J. F., E. BARBA-MACÍAS, C. TRINIDAD-OCAÑA, J. JUÁREZ-FLORES. 2016. Diversidad de crustáceos en la cuenca baja del río Papaloapan, Veracruz, México. *Hidrobiológica* 26 (3): 475-482.
- MIRES, D. 1983. The development of freshwater prawn (*Macrobrachium rosenbergii*) culture in Israel. *Bamidgeh* 3 (35): 63-72.
- NOM-127-SSA1-1994, "Salud ambiental, agua para uso y consumo humano-límites permisibles de calidad y tratamientos a que debe someterse el agua para su potabilización". 1994. <http://www.salud.gob.mx/unidades/cdi/nom/127ssa14.html> Reviewed: Jan 2, 2024.
- NOM-001-SEMARNAT-2021. Que establece los límites permisibles de contaminantes en las descargas de aguas residuales en cuerpos receptores propiedad de la nación. Available online at: [[https://www.dof.gob.mx/nota\\_detalle.php?codigo=5645374&fecha=11/03/2022#gsc.tab=0](https://www.dof.gob.mx/nota_detalle.php?codigo=5645374&fecha=11/03/2022#gsc.tab=0)]. (download January 2, 2024).
- PRETTO, M. R. 1988. *Manual del cultivo del camarón de río gigante de Malasia. Dirección Nacional de acuicultura rural*. CEDIA. Santiago de Varaguas. Panamá 122 p.
- SALAS-MONREAL, D., M. L. RIVERÓN-ENZASTIGA, J. J. SALAS-PÉREZ, R. BERNAL-RAMÍREZ, M. MARÍN-HERNÁNDEZ & A. GRANADOS-BARBA. 2019. Bathymetric flow rectification in a tropical micro-tidal estuary. *Estuarine, Coastal and Shelf Science* 235, 106562 DOI: 10.1016/j.ecss.2019.106562
- SALAS-MONREAL, D., V. DÍAZ-HERNÁNDEZ, J. A. ÁKE-CASTILLO, A. GRANADOS-BARBA & M. L. RIVERÓN-ENZASTIGA. 2020. Variación anual de los parámetros hidrográficos en la confluencia del río Jamapa y arroyo Moreno (México). *Intropica* 15 (1):59-65. DOI: 10.21676/23897864.3402
- SECRETARÍA DE MEDIO AMBIENTE Y RECURSOS NATURALES (SEMARNAT). 2002. Comisión Nacional del Agua, Gerencia de Saneamiento y Calidad del Agua. Calidad del agua del Río Jamapa 1 conforme a parámetros físicos, Cuadro III.2.2.16 químicos y biológicos, 1990-2001. Estación de medición: El Tejar, Veracruz (96°10'13»long. O; 19°03'29»lat. N). Available online at: [[http://www.paot.org.mx/centro/inesemarnat/informe02/estadisticas\\_2000/compendio\\_2000/03dim\\_ambiental/03\\_02\\_Agua/data\\_agua/Cuadrol-III.2.2.16.htm](http://www.paot.org.mx/centro/inesemarnat/informe02/estadisticas_2000/compendio_2000/03dim_ambiental/03_02_Agua/data_agua/Cuadrol-III.2.2.16.htm)]. (downloaded February 17, 2024).
- SOKAL, R. R. & F. J. ROHLF. 1995. *Biometry: the principles and practice of statistics in biological research*. W.H. Freeman, New York. 778 p.
- URBANO, T. A., A. L. SILVA-MEDINA, C. MORENO, M. GUEVARA & C. GRAZIANI. 2010. Crecimiento del camarón de agua dulce *Macrobrachium jelskii* (Miers, 1877), en lagunas de cultivo. *Zootecnia Tropical* 28 (2): 163-171.
- WILLIAMS, A. B. 1984. *Shrimps, lobsters and crabs of the Atlantic coast of the eastern United States, Maine to Florida*. Smithsonian Institution Press, Washington, D. C. 550 p.
- WoRMS. EDITORIAL BOARD. 2024. World register of marine species. Cariidea. Available online at: <http://marinespecies.org/aphia.php?p=taxdetails&id=106674> (downloaded on February 09 2023)
- ZAVALA-HIDALGO, J., SALMERÓN, O., AGUILAR, V., CERDEIRA, S. & M. KOLB. 2006. Caracterización y regionalización de los procesos oceanográficos de los mares mexicanos. CONABIO, Ciudad de México. Available online at: [[http://www.conabio.gob.mx/gap/index.php/Procesos\\_oceanogr%C3%A1ficos](http://www.conabio.gob.mx/gap/index.php/Procesos_oceanogr%C3%A1ficos)]. (downloaded on January 2, 2024).
- ZUUR, A. F., E. N. IENO & G. M. SMITH. 2007. *Analysing ecological data*. Springer, New York, London 370 p.