

## Macrobenthic Fauna of the Meric River (Turkish Thrace): Composition of the community as related to water quality

## Fauna macrobentónica del río Meric (Tracia turca): composición de la comunidad en relación a la calidad del agua

Menekse Tas-Divrik<sup>1\*</sup> and Timur Kirgiz<sup>2</sup>

Recibido: 12 de septiembre de 2017.

Aceptado: 13 de noviembre de 2018.

Publicado: 14 de diciembre de 2018.

### ABSTRACT

**Background:** Benthic macroinvertebrates are commonly used as indicators of the biological condition of waterbodies. Both the benthic macroinvertebrates and the physicochemical properties of the water provide important information about the pollution of a water environment. **Goals:** To investigate the relationships between assemblage composition of Oligochaeta (Annelida), Chironomidae (Diptera), and other macroinvertebrates with physical and chemical water variables in the Meric River, Turkey. **Methods:** This study was carried out by taking water and benthic samples at 8 stations at monthly intervals in order to determine what macrobenthic fauna exist in the Meric River and what environmental properties affect their distribution. **Results:** A total of 39 taxa were found and we determined that there is an average of 851 individuals per m<sup>2</sup>. Also, this study established the first records for *Brachiura sowerbyii* (Oligochaeta) and *Pottashia alternis* (Chironomidae) in the Turkish Thrace region. According to Shannon-Wiener index, while it was determined that Meric River has the highest diversity values with  $H' = 0.845$  at the station 4, the river has the poorest diversity with  $H' = 0.477$  at the station 2. Also, we examined similarities of distribution of Oligochaeta taxa by station and month using the Bray-Curtis index. Accordingly, while stations 6 and 8 were determined to be the most similar to each other, stations 2 and 5 were the least similar. The relationships between Oligochaeta taxa and physical and chemical parameters of water were evaluated using the Spearman Correlation index. As a result, we found that water temperature, pH, electrical conductivity, hydrogen sulfide, calcium, magnesium, phosphate, chlorine, salinity, dissolved oxygen and biological oxygen have positive correlations with some species of Oligochaeta, while nitrate and nitrite have negative correlations with some species of Oligochaeta. **Conclusions:** We made a number of suggestions for sustainable usage of this river.

<sup>1</sup> Sarkisla Asik Veyssel Vocational School, Cumhuriyet University, Sarkisla, 58400, Turkey.

<sup>2</sup> Department of Biology, Faculty of Science, Trakya University, Edirne, 22030, Turkey

\*Author for correspondence:  
menekse.tas@cumhuriyet.edu.tr

**Keywords:** Chironomidae, fauna, macrobenthic, Meric River, Oligochaeta

### RESUMEN

#### To cite as:

Tas-Divrik M. and T. Kirgiz. 2018. Macrobenthic Fauna of the Meric River (Turkish Thrace): Composition of the community as related to water quality. *Hidrobiológica* 28 (3): 277-285.

DOI:10.24275/uam/izt/dcbshidro/2018v28n3/Tas

**Antecedentes:** Los macroinvertebrados bentónicos se usan comúnmente como indicadores de la condición biológica de los cuerpos de agua. Estos organismos al igual que las propiedades fisicoquímicas del agua, proporcionan información importante sobre la contaminación de un ambiente acuático. **Objetivos:** Investigar las relaciones entre la composición del ensamblaje de Oligochaeta (Annelida), Chironomidae (Diptera) y otros macroinvertebrados con las variables físicas y químicas del agua en el río Meric, Turquía. **Métodos:** Se estudiaron muestras de agua y de bentos, tomadas en 8 estaciones en intervalos mensuales para determinar la fauna macrobentónica del río Meric y las propiedades ambientales que afectan su distribución. **Resultados:** Se encontraron un total de 39 taxa, con un promedio de 851 individuos por m<sup>2</sup>. Además las especies: *Brachiura sowerbyii* (Oligochaeta) y *Pottashia alternis* (Chironomidae), constituyen los primeros registros en la región de la Tracia turca. Según el índice de Shannon-Wiener, se determinó que el río Meric tiene los valores más altos de diversidad ( $H' = 0.845$ ) en la estación 4, y los más bajos ( $H' = 0.477$ ), en la estación 2. Además, se examinaron las similitudes de distribución de los taxones de Oligochaeta para las estaciones y los meses utilizando el índice de Bray-Curtis. Las estaciones 6 y 8 se determinaron como las más similares entre sí, por el contrario las estaciones 2 y 5 como las más diferentes. Las relaciones entre los taxones de Oligochaeta y los parámetros físicos y químicos del agua se evaluaron utilizando el índice de correlación de Spearman. Como resultado, se observó que la temperatura del agua, el pH, la conductividad eléctrica, el sulfuro de hidrógeno, el calcio, el magnesio, el fosfato, el cloro, la salinidad, el oxígeno disuelto y el oxígeno biológico tienen correlaciones positivas con algunas especies de Oligochaeta, mientras que el nitrato y el nitrito tienen correlaciones negativas con los organismos de ese grupo. **Conclusiones:** Se hacen sugerencias para el uso sustentable del río.

**Palabras clave:** Chironomidae, fauna, macrobentónico, Oligochaeta, río Meric

## INTRODUCTION

The Maritza/Evros/Meric river basin, which includes the Arda, Tunca, and Ergene tributaries, is one of the major river systems located in the eastern Balkans, with a total length of 550 km and total catchment area of 53000 km<sup>2</sup>. About 66% belongs to Bulgaria, 28% to Turkey, and 6% to Greece. The Meric is an international river that runs through an area of approximately 321 km in Bulgarian territory and flows into Aegean Sea. The Basin includes its main tributaries, i.e., the Arda and Tunca Rivers that mainly flow in Bulgaria and the Ergene River that flows entirely in Turkish territory. The basin provides suitable conditions for intensive irrigation and raising livestock (UNECE, 2009).

The Meric River forms the border between Greece and Turkey. Turkish authorities consider it to be an “international river” between Turkey and Greece and a “transboundary river” where it crosses the border between Bulgaria and Turkey. The water is used mostly for irrigation. It is also used at hydroelectric power plants. In addition, the Delta region provides good fishing for Turkey and Greece. The Delta is a very important stopover for birds that winter there. The Meric Delta is listed as a Class A International Wetlands (TCSV, 1989).

Many studies of the Bulgarian portion of river have been undertaken by Russev (1970), Russev *et al.* (1981), Detcheva (1981), Detcheva (1982), Russev and Janeva (1983), Kovachev (1985), Georgiev (2006), Stefanova *et al.* (2008), Rozdina *et al.* (2008), the oligochaeta studies by Dimitrov (1966), Uzunov (1976), Uzunov (1980), Uzunov (1981), Uzunov and Kovachev (1981), Uzunov *et al.* (1981), Uzunov and Kapustina (1993). Up to now, Özkan (1998) and Kalebaşı (1994) have studied the Turkish portion of the Meric River. There has been no study on macrobenthic fauna in the Turkish part of Meric River. To date, there have been no studies related to macrobenthic fauna and environmental parameters.

The objective of this study is to investigate the relationships between the physicochemical parameters and macrobenthic invertebrates in the Meric River. In addition, as part of the conclusions we offer a number of suggestions for the sustainable use of the river.

## MATERIALS AND METHODS

The Meric River is the most important aquatic ecosystem in the Thrace Region of Turkey and the longest river in the Balkans. It originates in Bulgaria, delimits a small segment of border between Greece and Bulgaria, and then establishes the entire Greek-Turkish border. After crossing the Greek - Turkish border, one of its main tributaries, the Arda, flows into the Meric, and after crossing the Bulgarian-Turkish border, the Tunca, the other main tributary, joins the Meric near the Turkish city of Edirne. The river is between 150 and 300 meters wide (Emir, 1990). The transboundary basin of the Meric is shared by three European countries: Bulgaria, Greece, and Turkey. Bulgaria is upstream; Turkey and Greece are downstream (Fig. 1).

We conducted this study from January 2011 to December 2011 at eight stations. Samples were taken monthly. Some properties of selected stations on the river: Station 1: This station is located upstream from industrial facilities located in Kapıkule. The substrates are composed of mud. The river's width at this station is about 160 meters. Station 2: This station is about 5 km away from the station 1. The substrates are composed of sandy-mud. The river's width at this station is about 175 meters. Station 3: The Arda Stream joins the Meric. The substrates are

composed of sand. The width at this station is about 180 meters. Station 4: At this station, the Tunca Stream joins the Meric. The substrates are composed of clay-detritus. The width at this station is about 200 meters. Station 5: Tatarköy village. There are many rice-growing areas around this location. This station is used for irrigation. The substrates are composed of stones. The river's width at this station is about 220 meters. Station 6: Saclımüsellim village. This station is located 23 km from station 5. The substrates are composed of sand. The width at this station is about 200 meters. Station 7: Like station 5, this station has many rice fields. This station is located before the junction with the river Ergene. The substrates are composed of sand. The river's width at this station is about 210 meters. Station 8: This station is located after the point of union with the river Ergene. The substrates are composed of clay and stone. The width of this station is about 250 meters.

At each site, water samples were taken periodically, at which time samples of benthic macroinvertebrates were taken. Benthic macroinvertebrate samples were taken at a depth of 1.5 meters.

Macrobenthic samples were taken from each station twice by using Ekman Birge grab (15 x 15 cm) and washed through on sieve series (1.19 mm, 0.595 mm, 0.297 mm mesh size). All collected samples were immediately fixed in 4% formaldehyde in the field and then transferred to 70% ethanol. In the laboratory, collected benthic macroinvertebrates were sorted and counted by using a stereomicroscope and then identified to the lowest possible taxon (species, genus, or families) by using a binocular microscope.

Identification of the Oligochaeta species was made by using keys in Brinkhurst and Jamieson (1971), Brinkhurst and Wetzel (1984), Kathman and Brinkhurst (1998), Milligan (1997), Timm (1999), and Wetzel *et al.* (2000). Chironomidae larvae were identified by using the keys in Saether (1980), Cranston (1982), Pinder and Reiss (1983).

At each station, the Meric River's water temperature (using ordinary thermometer), electrical conductivity (using a conductivity meter), and pH (using a pH meter) were measured when benthic samples were taken. The latter were taken by a Ruttner sampler and carried to the laboratory in 2 L bottles with dissolved oxygen (using the classical Winkler method). BOD, SO<sub>4</sub><sup>2-</sup>, PO<sub>4</sub><sup>2-</sup>, NO<sup>3</sup>-N, NO<sup>2</sup>-N, Cl, Mg, Ca, total hardness, salinity, H<sub>2</sub>S, and suspended solid material were analyzed by classical titrimetric and spectrophotometric methods. The water-quality level was determined according to official standards set by Turkey's National Water Quality Standards for inland waters (SKKY, 2004). The Shannon-Wiener index was used to evaluate the species diversity of the river. The similarities between the stations and months were evaluated by the Bray-Curtis similarity index (Krebs, 1999). The relationships between Oligochaeta taxa and physicochemical parameters of the water were evaluated by using the Spearman Correlation index (Krebs, 1999).

## RESULTS

A total of 39 taxa consisting of an average of 851 individuals/m<sup>2</sup> were collected at the Meric River stations during the study period. Samples were grouped as “Oligochaeta”, “Chironomidae” and “other benthic macroinvertebrates”. We found that 13 taxa belonging to Oligochaeta consist of 686 individuals/m<sup>2</sup>, 17 taxa belonging to Chironomidae consist of 145 individuals/m<sup>2</sup>, and 9 taxa belonging to other benthic macroinvertebrates consist of 20 individuals/m<sup>2</sup>. *Brachiura sowerbyi* (Beddard, 1982) that belongs to Oligochaeta and *Pottashia alternis* (Sa-




Figure 1. Map of the Meric River (Turkish Thrace) and sampling stations.

hin, 1987) that belongs to Chironomidae were determined as new records for Turkish fauna. With regard to the rational distribution of these groups, Oligochaeta were the predominant group with 81%, followed by Chironomidae larvae with 17%, and other benthic macroinvertebrates 2% (Table 1).

The most abundant taxa on average was *Limnodrilus hoffmeisteri* (Claparède, 1862), comprising 69.24%, while the least abundant taxa on average were *Limnodrilus profundicola* (Verrill, 1871), *Nais bretscheri*, (Michaelson, 1899), *Aulophorus furcatus* (Müller, 1774), *Lumbriulus* sp., and Enchytraeidae 0.14%. In addition, *L. hoffmeisteri* is the most abundant species at station 5. We found that the abundance of Oligochaeta was very high at station 5 (total number of individuals was 1941 per m<sup>2</sup>), while station 2 had the least (total number of individuals was 135 per m<sup>2</sup>) (Table 1).

Oligochaeta were found to have the highest number in May. In February and October none were found. The number of taxa belonging to Oligochaeta was the highest in June (9 taxa) (Table 2). Oligochaetes were found to have the highest number of taxa at station 4 (7 taxa), while the least taxa were found at station 2 (3 taxa) (Table 1).

Chironomidae had 17 taxa and 17 % abundance. *Chironomus Camptochironomus tentans* (Fabricius, 1805) was found to have the highest abundance (40%), while *P. (Holotanypus) sp.*, *Rheocricotopus fuscipes* (Kieffer, 1909), *Stictrochironomus sp.*, *Pentapedilum executum* (Kieffer, 1915), *Paratendipes albimanus* (Meigen, 1818), *Einfeldia pagana* (Meigen, 1838), *Potthastia alternis* (Sahin, 1987), and *Rheo-*

*tanytarsus* sp., were found to have the lowest abundance (0.68%). Chironomids were found to have the highest number of taxa at station 4 (12 taxa), followed by station 7 (9 taxa), and station 1 (8 taxa) respectively. *C. tentans*, *Chironomus anthracinus* (Zetterstedt, 1860) and *Cryptochironomus defectus* (Kieffer, 1913) were the most abundant and frequently recorded on sediments. Chironomidae was found to have the highest number in April. In January and February, none were found (Table 2). The number of taxa belonging to Chironomidae was the highest in May (11 taxa) (Table 2).

Other benthic macroinvertebrates had 9 taxa and 2% abundance. Odonata belonging to this group was found to have the highest abundance (30%), while Diptera (Chironomidae larvae excepted), Bivalvia, Coleoptera larvae, Trichoptera, Hemiptera, and Isopoda were found to have the lowest abundance (5%). Other benthic macroinvertebrates were found to have the highest number at stations 4 and 5 (total number of individuals 45 per m<sup>2</sup>), following by station 1 (29 per m<sup>2</sup>), and station 7 (15 per m<sup>2</sup>). None were found at station 6 (Table 1). Other benthic macroinvertebrates were found to have the highest number in April (483 per m<sup>2</sup>), following by June (433 per m<sup>2</sup>), and July (308 per m<sup>2</sup>), respectively. In January and February, none were found. The number of taxa belonging to other benthic macroinvertebrates was highest in May (6 taxa) (Table 2).

According to the Shannon-Weiner index, the Meric River had H' = 0.69 richness on average and station 4 was found to have the highest diversity H' = 0.84, while station 2 had the lowest H' = 0.47.

Table 1. List of taxa and number of individuals (per square meter) of benthic macroinvertebrates that were recorded at stations along the Meric River (Turkish Thrace). Ave. = Average; Ab. = Abundance; Tot.Ab.= Total abundance.

Station/Benthic macroinvertebrates	1	2	3	4	5	6	7	8	Ave.	Ab.	Tot.Ab.
<b>Oligochaeta</b>											
<i>Tubifex tubifex</i> (Müller,1774)	0	9	11	13	402	4	20	15	59	8.60	6.93
<i>Limnodrilus hoffmeisteri</i> (Claparède,1862)	819	124	176	691	976	346	257	413	475	69.24	55.81
<i>L. dekemianus</i> (Claparède,1862)	44	2	2	54	0	0	0	18	15	2.19	1.76
<i>L. profundicola</i> (Verrill,1871)	0	0	0	0	0	0	2	0	1	0.14	0.11
<i>Potamothrix hammoniensis</i> (Michaelsen,1901)	11	0	2	40	239	6	2	29	41	5.98	4.81
<i>Nais bretscheri</i> (Michaelsen,1899)	2	0	0	0	0	0	0	0	1	0.14	0.11
<i>N. elinguis</i> (Müller,1773)	0	0	15	179	0	24	200	0	52	7.58	6.11
<i>Aulophorus furcatus</i> (Müller, 1774)	0	0	0	2	0	0	0	0	1	0.14	0.11
<i>Dero digitata</i> (Müller, 1773)	0	0	0	0	259	0	0	4	32	4.67	3.76
<i>D. obtusa</i> (D'Ukem,1855)	0	0	0	0	26	0	0	0	3	0.44	0.35
<i>Ophidonais serpentina</i> (Müller, 1773)	0	0	0	0	39	0	0	0	4	0.60	0.47
<i>Lumbriculus</i> sp.	0	0	0	2	0	0	0	0	1	0.14	0.11
Enchytraeidae	2	0	0	0	0	0	0	4	1	0.14	0.11
Total Oligochaeta	878	135	206	981	1941	380	481	483	686	100	81
Number of taxa	5	3	5	7	6	4	5	6			
<b>Chironomidae</b>											
<i>Tanytus punctipennis</i> (Meigen, 1818)	0	0	2	2	50	3	0	0	7	4.82	0.82
<i>Praccladius</i> ( <i>Holotanytus</i> ) sp.	0	0	0	3	8	0	0	0	1	0.68	0.11
<i>Cricotopus</i> ( <i>C.</i> ) <i>bicinctus</i> (Meigen, 1818)	2	0	0	9	0	0	5	0	2	1.40	0.23
<i>Rheocricotopus fuscipes</i> (Kieffer, 1909)	2	0	0	0	0	0	0	0	1	0.68	0.11
<i>Chironomus anthracinus</i> (Zetterstedt, 1860)	2	0	11	11	48	7	4	0	10	6.90	1.17
<i>C. (C.) tentans</i> (Fabricius, 1805)	3	1	11	4	365	57	0	28	58	40	6.81
<i>C. plumosus</i> (Linnaeus,1758)	2	0	0	5	28	6	0	0	5	3.45	0.58
<i>Stictochironomus</i> sp.	0	0	0	2	0	0	5	0	1	0.68	0.11
<i>Polypedilum aberrans</i> (Chernovskij,1949)	26	0	27	153	116	16	18	0	44	30.34	5.17
<i>P. exsectum</i> (Kieffer,1915)	0	0	0	0	0	0	2	0	1	0.68	0.11
<i>Paratendipes albimanus</i> (Meigen, 1818)	0	0	0	0	0	0	4	0	1	0.68	0.11
<i>Cryptochironomus defectus</i> (Kieffer, 1913)	12	2	8	18	0	0	2	0	5	3.45	0.60
<i>Einfeldia pagana</i> (Meigen, 1838)	0	0	0	0	0	0	0	3	1	0.68	0.11
<i>Pottashia alternis</i> (Sahin, 1987)	0	0	0	0	0	2	2	0	1	0.68	0.11
<i>Tanytarsus gregarius</i> (Kieffer, 1909)	3	0	0	24	5	0	2	2	4	2.8	0.47
<i>Virgotanytarsus arduensis</i> (Kieffer, 1909)	0	0	0	22	0	0	0	0	2	1.40	0.23
<i>Rheotanytarsus</i> sp.	0	0	0	2	0	0	0	0	1	0.68	0.11
Total Chironomidae	52	3	59	255	620	91	44	33	145	100	17
Number of taxa	8	2	5	12	7	6	9	3			
<b>Other benthic macroinvertebrates</b>											
Diptera	0	0	2	2	2	0	0	0	1	5	0.1
Gastropoda	0	0	0	0	33	0	0	7	5	25	0.5
Bivalvia	0	0	0	0	0	0	9	0	1	5	0.1
Ephemeroptera	4	2	0	15	0	0	0	0	3	15	0.3
Odonata	18	2	2	19	6	0	2	4	6	30	0.7
Coleoptera (larvae)	2	0	0	0	0	0	0	0	1	5	0.1
Trichoptera	5	0	0	2	2	0	0	0	1	5	0.1
Hemiptera	0	0	0	7	2	0	0	0	1	5	0.1
Isopoda	0	0	0	0	0	0	4	0	1	5	0.1
Total other benthic macroinvertebrates	29	4	4	45	45	0	15	11	20	100	2
Number of taxa	4	2	2	5	5	0	2	2			



The results of Bray-Curtis index indicated that stations 6 and 8, stations 1 and 4, and stations 2 and 3 are the most similar to each other (82.50%, 80.25%, and 79.17% similarities, respectively). Stations 2 and 5, stations 5 and 7, stations 1 and 2 were determined to be the least similar (12.81%, 23.87%, 24.87% similarities, respectively). The Bray Curtis Similarity index indicated that September and July, July and November, September and November are the most similar to each other (92.15%, 89.56%, 85.79% similarities, respectively). February and April, February and June were determined to be the least similar (0.07%, 0.15% similarities, respectively) (Figs 2a-b).


The average data on physicochemical parameters at each station are shown in Table 3. The lowest electrical conductivity was 214  $\mu\text{s}/\text{cm}$  (at station 8), while the highest was 1197  $\mu\text{s}/\text{cm}$  (station 8). The average value measured was 502  $\mu\text{s}/\text{cm}$ . The lowest pH measured was 7.50 (station 8), while the highest was 10.03 (station 1). The average value was 8.42. The lowest dissolved oxygen was 2.28 mg/L (station 2), while the highest was 8.94 (station 7). The average value of dissolved oxygen was 5.53 mg/L. The values of biological oxygen demand ranged between 6.75-78.7 mg/L. The average value of biological oxygen demand was 37.32 mg/L. The lowest sulphate registered was 0.641 mg/L (station 4), while the highest was 4.250 mg/L (station 8). The average value of sulphate was 1.841 mg/L. The lowest phosphate value was 0 mg/L (stations 3, 4, 6, and 7), while the highest was 0.273 mg/L (station 5). The average phosphate value was 0.068 mg/L. The nitrite values ranged between 0-0.641 mg/L. The average value of nitrite was 0.034 mg/L. The nitrate values ranged between 0-33.750 mg/L. The average nitrate value 5.846 mg/L. The lowest chloride was 3.99 mg/L (station 1), while the highest was 179.94 mg/L (station 8). The average chloride value was 34.99 mg/L. The lowest magnesium value was 0.96 mg/L (station 4), while the highest was 59.56 mg/L (station 6). The average magnesium value was 16.71 mg/L. The lowest calcium was 24.04 mg/L (station 5), while the highest was 90.58 mg/L (station 8). The average calcium value was 54.75 mg/L. The lowest value of total hardness was 3.2 FS° (station 4), while the highest was 41.2 (station 6). The average total hardness was 17.3 FS°. The lowest salinity was 0.02‰ (station 4), while the highest was 0.23 (station 8). The average value was 0.068‰. Hydrogen sulfide values ranged between 0-3.195 mg/L. The average value of hydrogen sulfide was 0.110 mg/L. Suspend solid material registered ranged between 70-1630 mg/L. The average value of suspend solid material was 345 mg/L (Table 3).

**DISCUSSION**

Oligochaeta was the dominant group of benthos in the river, while Chironomidae was the second dominant group of benthic fauna during the study period. Oligochaeta and Chironomidae larvae are usually abundant in benthos and play important roles in river ecosystems. The main reasons are that oligochaetes also feed on organic material in the water and are common in environments where large amounts of organic materials are present due to their ability to survive with low oxygen levels and other low-level conditions better than most other macrobenthos species (Yankson & Kendall, 2001). Like oligochaetes, chironomid larvae have also been used as indicators of organic pollution because they are often abundant in environments with low oxygen where organic material as a food resource is abundant (Coffman & Ferrington, 1996; Jenderedjian *et al.*, 2007). The composition and distribution of Oligochaeta species depend on many factors such as water temperature,

physical and chemical properties of the water, sediments, microfauna, and vegetation (Grigelis *et al.*, 1981). Up to now, there has been no study of the Meric River regarding benthic fauna and their relationships with physicochemical parameters. For this reason, there are no other studies to compare with our work. No Oligochaetes were found in February and October. In February, rain and snow may have increased the water level in the river. Thus, sampling may not have been done exactly at the river bed. In October, snowfall caused the water level in the river to rise and sampling could not be done.

In Özkan's study (1998), the average number of Chironomidae larvae was 481 per m<sup>2</sup> for 65 taxa. In our study, we found that the average number of Chironomidae larvae was 145 per m<sup>2</sup> for 17 taxa. If this study is compared with Özkan's findings (1998), we see that both the individual and taxa number of Chironomidae larvae are declining. The main reasons for the decline of this species are habitat loss, pollution, meteorological factors, and changes in the structure of the river bottom.



Figures 2a-b. Bray Curtis Similarity Dendrograms by sampling stations (a) and months (b) along the Meric River (Turkish Thrace) based on Oligochaeta species. st = station.

Table 2. Monthly distribution of benthic macroinvertebrates and number of individuals (per square meter) in the Meric River (Turkish Thrace).

Month/Benthic macroinvertebrates	Jan.	Feb.	Mar.	Apr.	May.	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
<b>Oligochaeta</b>												
<i>Tubifex tubifex</i> (Müller, 1774)	0	0	0	0	278	330	5	0	8	0	31	59
<i>Limnodrilus hoffmeisteri</i> (Claparède, 1862)	0	0	97	481	2038	889	319	508	353	0	291	722
<i>L. udekemianus</i> (Claparède, 1862)	0	0	0	0	0	0	0	86	5	0	0	22
<i>L. profundicola</i> (Verrill, 1871)	0	0	0	0	69	3	0	0	0	0	0	0
<i>Potamothrix hammoniensis</i> (Michaelson, 1901)	0	0	11	155	295	11	15	4	5	0	0	0
<i>Nais bretscheri</i> (Michaelson, 1899)	0	0	0	0	0	3	0	0	0	0	0	0
<i>N. elinguis</i> (Müller, 1773)	0	0	109	492	22	5	0	0	0	0	0	0
<i>Aulophorus furcatus</i> (Müller, 1774)	0	0	0	0	0	0	0	0	4	0	0	0
<i>Dero digitata</i> (Müller, 1773)	0	0	0	308	81	0	0	5	0	0	0	0
<i>D. obtusa</i> (D'Ukern, 1855)	0	0	0	0	0	39	0	0	0	0	0	0
<i>Ophidonais serpentina</i> (Müller, 1773)	0	0	0	58	0	0	0	0	0	0	0	0
<i>Lumbriculus</i> sp.	0	0	0	0	0	3	0	0	0	0	0	0
<i>Enchytraeidae</i>	5	0	0	0	0	3	0	0	0	0	0	0
Total	5	0	217	1494	2783	1286	339	603	375	0	322	803
Number of taxa	1	0	3	5	6	9	3	4	5	0	2	3
<b>Chironomidae</b>												
<i>Tanytus punctipennis</i> (Meigen, 1818)	0	0	0	0	3	75	0	3	3	0	0	0
<i>Pracladius (Holotanytus)</i> sp.	0	0	0	0	0	11	0	3	0	0	0	0
<i>Cricotopus (C.) bicinctus</i> (Meigen, 1818)	0	0	5	11	5	0	0	0	0	0	0	3
<i>Rheocricotopus fuscipes</i> (Kieffer, 1909)	0	0	0	0	0	0	3	0	0	0	0	0
<i>Chironomus anthracinus</i> (Zetterstedt, 1860)	0	0	0	22	17	66	8	8	3	0	0	0
<i>C. (C.) tentans</i> (Fabricius, 1805)	0	0	0	308	230	94	30	8	8	0	13	8
<i>C. plumosus</i> (Linnaeus, 1758)	0	0	0	30	19	0	3	8	0	0	0	0
<i>Stictochironomus</i> sp.	0	0	0	0	3	0	0	7	0	0	0	0
<i>Polypedilum aberrans</i> (Chernovskij, 1949)	0	0	0	94	14	129	200	30	69	3	0	0
<i>Pentapedilum exectum</i> (Kieffer, 1915)	0	0	0	0	3	0	0	0	0	0	0	0
<i>Paratendipes albimanus</i> (Meigen, 1818)	0	0	0	6	0	0	0	0	0	0	0	0
<i>Cryptochironomus defectus</i> (Kieffer, 1913)	0	0	0	0	3	5	36	5	11	0	0	3
<i>Einfeldia pagana</i> (Meigen, 1838)	0	0	0	0	0	0	0	0	7	0	0	0
<i>Potthastia alternis</i> (Sahin, 1987)	0	0	0	3	3	0	0	0	0	0	0	0
<i>Tanytarsus gregarius</i> (Kieffer, 1909)	0	0	0	3	8	11	16	0	14	3	0	0
<i>Virgotanytarsus arduensis</i> (Kieffer, 1909)	0	0	0	0	0	0	0	0	33	0	0	0
<i>Rheotanytus</i> sp.	0	0	0	0	0	0	0	0	5	0	0	0
Total	0	0	5	477	308	391	296	72	153	6	13	14
Number of taxa	0	0	1	8	11	7	7	8	9	2	1	3
<b>Other benthic invertebrates</b>												
Diptera	0	0	0	0	10	3	0	0	0	0	0	0
Gastropoda	0	0	0	0	28	25	0	5	0	0	0	0
Bivalvia	0	0	0	0	3	0	3	0	0	0	0	0
Ephemeroptera	0	0	0	0	3	2	3	0	17	0	0	0
Odonata	0	0	0	3	3	7	6	3	36	7	6	5
Coleoptera (larvae)	0	0	0	0	0	5	0	0	0	0	0	0
Trichoptera (larvae)	0	0	0	3	3	0	0	5	0	3	0	5
Hemiptera	0	0	0	0	0	0	0	0	7	0	0	0
Isopoda	0	0	0	0	0	0	0	0	0	0	0	8
Total	0	0	5	483	358	433	308	88	213	19	19	32
Number of taxa	0	0	1	2	6	5	3	3	3	2	1	3

Uzunov (1980) reported that there were 8 families for a total of 79 species along the Bulgarian segment of the Meric River. In the study of the Meric River by Uzunov *et al.* (1981), the following taxa were found: Oligochaeta (54), Gastropoda (7), Bivalvia (2), Isopoda (1), Ephemeroptera (45), Odonata (9), Heteroptera (10), Diptera (39). In Russev and Jagneva's study (1983), 46 Ephemeroptera species were found. In Uzunov and Kapustina's study (1993), they found 54 Oligochaeta taxa.

According to SKKY (2004), the water temperature, pH, chloride, sulphate, phosphate, and nitrate values were found to be of first quality level. The value of dissolved oxygen was found to be at a second quality, whereas nitrite-N was found to be between second and third quality level. Biological oxygen demand was at fourth quality level. The relationships between the Oligochaeta species and the physicochemical parameters were evaluated by the Spearman correlation index. According to this index, the abundance of *Limnodrilus hoffmeisteri* (Claparède, 1862) showed positive correlations of water temperature ( $r = 0.627, p < 0.05$ ), electrical conductivity ( $r = 0.605, p < 0.05$ ), magnesium ( $r = 0.805, p < 0.01$ ), salinity ( $r = 0.717, p < 0.01$ ), chloride ( $r = 0.580, p < 0.05$ ), phosphate ( $r = 0.609, p < 0.05$ ), pH ( $r = 0.596, p < 0.05$ ). There was a negative correlation to nitrate ( $r = -0.600, p < 0.05$ ). The abundance of *L. udekemianus* showed positive correlations to hydrogen sulfide ( $r = 0.739, p < 0.01$ ), dissolved oxygen ( $r = 0.736, p < 0.01$ ), phosphate ( $r = 0.603, p < 0.05$ ), and biological oxygen demand ( $r = 0.709, p < 0.01$ ). The abundance of *Nais elinguis* (Müller, 1773) showed positive correlations to hydrogen sulfide ( $r = 1.000, p < 0.01$ ), and negative correlations to nitrate ( $r = -0.604, p < 0.05$ ). The abundance of *Tubifex tubifex* (Müller, 1774) showed positive correlations to calcium ( $r = 0.633, p < 0.05$ ), chloride ( $r = 0.640, p < 0.05$ ), and phosphate ( $r = 0.591, p < 0.05$ ). The abundance of *Potamothrix hammoniensis* (Michaelsen, 1901) showed

positive correlations to water temperature ( $r = 0.697, p < 0.05$ ), and negative correlations to nitrate ( $r = -0.636, p < 0.05$ ).

Compared with previous studies of Kalebaşı (1994) and Ozkan (1998), the river water has become alkaline. Dissolved oxygen content had dropped.

Electrical conductivity is higher at station 8 than at the other stations. The Ergene River has been adversely affected by increasing population, industrial activities, heavy pesticides and fertilizer use in agriculture, and domestic wastes. Water hardness was found to be at an intermediate level. Since there are no previous studies of water hardness, we cannot make a comparison. Suspended solid material averaged 348 mg/L in the river. Suspended solids in a body of water are often due to natural causes. These natural solids include organic and inorganic materials such as silt and sediment. The majority of suspended sediment in water bodies comes from runoff and erosion. Some of the more common suspended solid pollutants are wastewater effluent, sewage, and airborne particulates.

Rivers are open, dynamic ecosystems whose physical, chemical, and biotic characteristics are greatly influenced by anthropogenic activities within their drainage basins (Moyaka *et al.*, 2004).

The river, originating in Bulgaria, receives wastewater from industrial plants in the Kapıkule. At first, it merges with the Arda River, and later with the Tunca River. After passing through rice fields, it joins the Ergene River. The Ergene River has been adversely affected by increasing population, industrial activities, heavy pesticide and fertilizer use in agriculture, and domestic wastes. The Ergene River is the main source of pollution in the Meric Basin.

Table 3. The average physicochemical parameters of the Meric River (Turkish Thrace) during the study in 2011. W.T. = Water temperature; E.C. = Electrical Conductivity; D.O. = Dissolved oxygen; S.S.M. = Suspended Solid material; T.H. = Total hardness; BOD = Biological oxygen demand.

Months/ Parameters	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
W. T. (°C)	4.31	5.25	9.31	13.8	20.3	25.5	28.3	26.3	23.4	12	6.3	7.6
pH	8.86	8.17	7.96	8.13	8.4	9.08	8.48	8.41	8.34	8.16	8.26	8.45
E.C. (µS/cm)	295.6	349.7	411.2	417.1	539.1	619.8	649.2	618.3	685.3	453.7	510.1	484
D.O. (mg/L)	4.73	5.31	5.09	3.01	4.92	5.16	4.99	7.27	6.99	6.28	6.82	5.87
Salinity (‰)	0.06	0.06	0.06	0.05	0.06	0.09	0.06	0.06	0.08	0.07	0.08	0.07
Cl <sup>-</sup> (mg/L)	27.4	27.8	33.4	23.4	30.9	35.2	42.3	30.8	53.9	32.3	41.1	40.9
H <sub>2</sub> S (mg/L)	0.000	0.000	0.026	0.346	0.266	0.399	0.000	0.213	0.079	0.000	0.000	0.000
S.S.M. (mg/L)	383.7	270	360	343.7	372.5	448.7	462.5	200	342.5	287.5	373.7	347.5
Ca <sup>+2</sup> (mg/L)	44.28	43.37	47.19	55.5	56.7	58.01	41.87	57.51	63.82	58.21	71.63	59.71
Mg <sup>+2</sup> (mg/L)	7.07	7.62	8.77	14.52	12.4	14.76	24.57	17	24.51	16.46	28.6	22.88
T. H. (FS°)	11.2	13.7	15.4	7.85	9.02	8.37	20.6	20.67	26.05	21.07	29.7	24.92
NO <sup>2</sup> -N (mg/L)	0.03	0.028	0.053	0.02	0.019	0.08	0.024	0.016	0.074	0.028	0.028	0.007
NO <sup>3</sup> -N (mg/L)	8.318	8.191	9.643	6.258	5.041	3.426	0.683	2.759	7.751	3.504	4.153	10.44
PO <sub>4</sub> <sup>-3</sup> (mg/L)	0.041	0.041	0.075	0.053	0.061	0.044	0.02	0.064	0.085	0.052	0.069	0.227
SO <sub>4</sub> <sup>-2</sup> (mg/L)	1.044	1.218	1.567	1.037	2.07	1.908	2.383	1.798	2.242	2.045	2.873	1.952
BOD (mg/L)	15.83	19.47	21.2	11.46	21.66	40.41	40.55	62.85	61.77	53.96	54.25	44.65

The river, an important irrigation source for the Thrace region, is a natural resource that must be used and preserved in a stable manner. As a result, both physicochemical properties and the results of tests on macrobenthic fauna indicate that the water quality of this river is organically polluted.

In recent years, due to population growth, rapid urbanization, and increasing discharge of solid wastes, water pollution problems are emerging.

Many industries still discharge untreated wastes and wastewaters into rivers. This should not be done if the goal is to assure sustainable use of the river water. Industrial wastewater contains pollutants. Only the necessary fertilizers and insecticides should be used at the appropriate dose. All the components that make up the river basin should be defined and managed together as a system. Because the Meric is both a trans-boundary and a border river, Turkey, Bulgaria, and Greece need to cooperate in order to develop and manage its waters. This would involve undertaking joint projects and studies. These and similar studies should be done regularly to monitor and determine how river water can be used sustainably to maintain its quality. The change of water quality in the river must be recorded.

### ACKNOWLEDGEMENTS

We would like to thank Dr. Belgin Camur Elipek and Dr. Burak Öterler for their helps during the field works and guidences. We would like to thank Dr. Nurten Hacet for Odonata diagnosis. This research has been supported by TUBAP 2010-168 Project.

### REFERENCES

- BRINKHURST, R. O. & B. G. M. JAMIESON. 1971. *Aquatic Oligochaeta of the World*. Oliver & Boyd, Edinburg.
- BRINKHURST, R. O. & M. J. WETZEL. 1984. *Aquatic Oligochaeta of the World: Supplement, A Catalogue of New Freshwater Species, Descriptions and Revisions*, No: 44, Canadian Technical Report of Hydrography and Ocean Sciences, Canada. 101 p.
- COFFMAN, W. & L. JR. FERRINGTON. 1996. Chironomidae. In: R. Merritt & K. Cummins (ed). *An Introduction to the Aquatic Insects of North America*. Dubuque, Iowa, USA, Kendall/Hunt Publishing Company. pp. 591-754.
- CRANSTON, P. S. 1982. A Key to the Larvae Orthocladinae of the British. *Freshwater Biological Association Scientific Publication* 4: 1-152.
- DETCHEVA, R. 1981. Etude de la composition et de la repartition saisoniere des Cilies de la riviere Maritza. *Hydrobiology* (Sofia) 14: 16-30.
- DETCHEVA, R. 1982. Caracteristiques ecologiques des Cilies de la riviere Maritza. *Annual Statistica Biologica Besse-en-Chandesse* 16: 200-219.
- DIMITROV, M. 1966. Chironomidae Larvae (Family Chironomidae, Diptera) and Oligochaeta (Order Oligochaeta) of the Maritsa River and Some of its Tributaries. In: Die Fauna Thrakiens (Sammelwerk), BAN, Zoologisches Institut und Museum, Band III, 309-318 (In Bulgarian, Russian summary).
- EMIR, B. D. 1990. Meric Basin Environmental Problem Research Project (Preliminary Report). Thrace University Environmental Problem Research and Application Center (In Turkish).
- GEORGIEV, D. G. 2006. The Fish Species Participation in Otters (*Lutra lutra*) Diet in Maritza River, West of Plovdiv Town (Southern Bulgaria). *Animalia* 42:153-159.
- GRIGELIS, A., R. LENKAITIS, O. NAINAITÉ & E. ŽUKAITĖ. 1981. Peculiarities of Distribution of Cold-Stenotherm Hydrobionts in Lakes of the National Park of the Lithuanian SSR. *Verhandlungen des Internationalen Verein Limnologie*, 21: 501-503.
- JENDEREDJIAN, K., S. HAKOBYAN & A. JENDEREDJIAN. 2007. Use of benthic invertebrates as indicators of pollution origin in agricultural and urban areas. In: A. Ebel, T. Divitashvili (ed). *Air, Water and Soil Quality Modelling for Risk and Impact Assessment*. NATO Security through Science Series C: Environmental Security, Netherlands: Springer Publishers. pp. 217-220.
- KALEBASI, Y. 1994. *Investigation of Chemical Pollution of Meric River*. Trakya University, Department of Chemistry, Master Thesis. (in Turkish with English abstract). 84 p.
- KATHMAN, R. D. & R. O. BRINKHURST. 1998. *Guide to The Freshwater Oligochaetes of North America*. Aquatic Resources Center, Tennessee, USA, 264 p.
- Kovachev, S. 1985. The Simuliid Complex of the Maritsa River as Example of a Highly Destructed Taxocenosis. *Hydrobiology* (Sofia) 26: 53-58 (In Bulg.).
- KREBS, C. J. 1999. *Ecological Methodology*. Addison Wesley Longman, Inc., Menlo Park, California. 620 p.
- MILLIGIAN, R. M. 1997. *Identification Manual for The Aquatic Oligochaeta of Florida Volume I, Freshwater Oligochaetes*. State of Florida Department of Environmental Protection Tallahassee, Florida, 187 p.
- MOYAKA, S. K., J. M. MATHOOKO & M. LEICHTFRIED. 2004. Influence of anthropogenic activities of water quality of a tropical stream ecosystem. *Africa Journal Ecology* 42: 281-288.
- OSRAM (MIDDLE EAST STRATEGIC RESEARCH CENTER). 2011. *Water Research Program*. "International Cooperation" Mandatory in Water Management of Meric River Basin, April 2011. Report No. 4: 1-47.
- OZKAN, N. 1998. The Dynamics of the Chironomidae (Diptera) Larvae in Some Tributary with Meric and Ergene Rivers, Trakya University, Institute of Natural Science, PhD Thesis, 143 p. (In Turkish with English abstract).
- PINDER, K. & F. RREISS. 1983. The Larvae of Chironominae (Diptera: Chironomidae) of the Holarctic Region- Keys and Diagnoses. *Entomologica Scandinavica Suppl.* 19: 293-435. Lund Sweden. ISSN:0105-3574.
- ROZDINA, D., G. RAIKOVA-PETEROVA, R. MARINOVA & E. UZUNOVA. 2008. Food Spectrum and Feding of *Barbus cyclolepis* Heckel from the Middle Stream of Maritza River (Bulgaria). *Bulgarian Journal of Agricultural Science* 14 (2): 209-213.
- RUSSEV, B. K., 1970. *Review of the Saprobiological Condition of the major Rivers in Bulgaria*. FAO. EIFAC, 70/SC III-2.



- RUSSEV, B. K. & I. J. JANEVA. 1983. The Significance of Mayflies (Ephemeroptera, Insecta) as Structural Constituents of Benthic Zoocenoses of the Maritsa River. *Hydrobiology* 19: 14-24.
- RUSSEV, B. K., J. I. UZUNOV, S. G. KOVACHEV, I. Y. YANEVA & L. S. IVANOVA. 1981. Tendencies of the Changes in the Saprobic Conditions of the Maritsa River. *Hydrobiology* 14: 51-64.
- SEATHER, O. A. 1980. Glossary of Chironomid Morphology Terminology (Diptera:Chironomidae). *Entomologica Scandinavica* Suppl. 14: 1-51. Lund, Sweden. ISSN:0105-3574.
- SKKY (SU KIRLIĞI KONTROL YÖNETMELİĞİ). 2004. *Water Pollution Control Regulation*. Published Official Journal. Date 31.12.2004, Number: 25687
- STEFANOVA, E., E. UZUNOVA, T. HUBENOVA, P. VASILEVA, D. TERZIYSKI & I. ILIEV. I. 2008. Age and Growth of Chub, *Leuciscus cephalus* L. from the Maritza River (South Bulgaria). *Bulgarian Journal of Agricultural Science* 14 (2): 214-220.
- TCSV (TURKEY ENVIRONMENTAL PROBLEMS FOUNDATION). 1989. *Wetlands of Turkey*. Publication of the Turkish Foundation for Environmental Problems.
- TIMM, T. 1999. *A Guide to the Estonian Annelida. Naturalist's Handbooks* 1. Estonian Academy Publishers, Tartu-Talinn. 208 p.
- UNECE (UNITED NATIONS ECONOMIC COMMISSION FOR EUROPE). 2009. *Convention on the Protection and Use of Transboundary Water Courses and International Lakes*. Transboundary Flood Risk Management: Experiences from the UNECE Region, United Nations Publication. ISBN:978-92-I-II70II-5.
- UZUNOV, J. 1976. Study on the water representatives of the class oligochaeta in Bulgaria. *Hydrobiology* (Sofia) 4: 71-75 (In Bulgarian).
- UZUNOV, J. 1980. Water Oligochets (Oligochaeta, Limicola) from Some Bulgarian Rivers. Frequency and Domination. *Hydrobiology* 12: 79-89.
- UZUNOV, J. 1981. Aquatic Oligochaetes (Oligochaeta) from the Maritsa River. *Hydrobiology* (Sofia) 15: 31-50 (In Bulgarian).
- UZUNOV, J., B. K. RUSSEV, S. G. KOVACHEV & I. J. JANEVA. 1981. Species List and Distribution of the Macrobenthos of the River Maritsa. *Hydrobiology* 14: 3-14.
- UZUNOV, J. & S. G. KOVACHEV. 1981. The Effect of the substrate on the Structure of the Macrozoobenthic Communities in the Maritsa River. *Hydrobiology* (Sofia) 15: 65-74 (In Bulgarian).
- UZUNOV, Y. & L. KAPUSTINA. 1993. Current Review on Oligochaeta from Macrozoobenthic Communities of the Bulgarian Rivers. *Lauterbornia* 13: 73-83.
- WETZEL, M. J., R. D. KATHMAN, S. V. FEND & K. A. COATES. 2000. *Taxonomy, Systematics and Ecology of Freshwater Oligochaeta*. Workbook Prepared for North American Benthological Society Technical Workshop, 48th Annual Meeting, Keystone Resort, CO. 120 pp. + app.
- YANKSON, K. & M. KENDALL. 2001. *A Student's Guide to the Seashore of West Africa*. Darwin Initiative Report 1, Ref. 162/7/45. 132.p